



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

 ScienceDirect

Applied Mathematics Letters 21 (2008) 780–785

**Applied  
Mathematics  
Letters**

[www.elsevier.com/locate/aml](http://www.elsevier.com/locate/aml)

# An analytical approximation to the solution of a wave equation by a variational iteration method

J. Biazar<sup>a,\*</sup>, H. Ghazvini<sup>a,b</sup>

<sup>a</sup> *Department of Mathematics, Faculty of Sciences, The University of Guilan, P.O. Box 1914, P.C. 41938, Rasht, Iran*

<sup>b</sup> *Department of Mathematics, School of Mathematical Sciences, Shahrood University of Technology, P.O. Box 316, P.C. 3619995161, Shahrood, Iran*

Received 7 June 2006; received in revised form 16 July 2007; accepted 3 August 2007

## Abstract

In this work, a variational iteration method, which is a well-known method for solving functional equations, has been employed to solve the general form of a wave equation which governs numerous scientific and engineering experimentations. Some special cases of wave equations are solved as examples to illustrate the capability and reliability of the method. The results reveal that the method is very effective. The restrictions of the method are mentioned.

© 2007 Elsevier Ltd. All rights reserved.

**Keywords:** Variational iteration method; Wave equation; Correction functional; Restricted variation; Lagrange multiplier

## 1. Introduction

Wave equations have attracted a great deal of attention and solving different forms of these equations has been an interesting task for mathematicians. Numerical methods which are commonly used such as finite difference and characteristics approaches need large amounts of computational work and usually the effect of rounding-off error causes loss of accuracy in the results. Analytical methods commonly used for solving wave equations are very restricted and can be used only in very special cases, so they cannot be used to solve equations for numerous realistic scenarios. The variational iteration method introduced by He [1–10] has been used by many mathematicians and engineers to solve various functional equations. It has been shown that this procedure is a powerful tool for solving various kinds of functional equations. For example, this scheme is used for solving delay differential equations [10]. Application of the variational iteration method to the Helmholtz equation is investigated in [11]. This technique is used for solving the Burgers and coupled Burgers equations [12]. Applications of the present method to coupled Schrödinger–KdV equations and the shallow water equation are provided [14]. The variational iteration technique is employed for solving an inverse problem [13]. The authors of this work also used this method for solving systems of ordinary differential equations, fourth-order parabolic equations and hyperbolic differential equations [15–17]. Also,

\* Corresponding author.

E-mail addresses: [biazar@guilan.ac.ir](mailto:biazar@guilan.ac.ir), [jbiazar@dal.ca](mailto:jbiazar@dal.ca) (J. Biazar).

this method was used to solve other nonlinear equations [18–22]. For the purpose of illustration of the methodology of the variational iteration method, let us consider the following nonlinear functional equation:

$$Lu(t) + Nu(t) = g(t)$$

where  $L$  is a linear operator,  $N$  is a nonlinear operator and  $g(t)$  is a known analytical function. According to the variational iteration method, we can construct the following correction functional:

$$u_{n+1}(t) = u_n(t) + \int_0^t \lambda(\xi)(Lu_n(\xi) + Nu_n(\xi) - g(\xi))d\xi,$$

where  $\lambda$  is a general Lagrange multiplier which can be identified via variational theory,  $u_0(t)$  is an initial approximation, with possible unknowns, and  $\tilde{u}_n$  is considered as a restricted variation [5,6], i.e.  $\delta\tilde{u}_n = 0$ . Therefore, we first determine the Lagrange multiplier  $\lambda$  that will be identified optimally via integration by parts. The successive approximations,  $u_{n+1}(t)$ , to the solution  $u(t)$  will be readily obtained upon using the Lagrange multiplier obtained and by using any selected function  $u_0$ . Consequently, the exact solution may be obtained by using  $u = \lim_{n \rightarrow \infty} u_n$ .

## 2. Solution of a wave equation by the variational iteration method

Consider the following general form of a wave equation:

$$\frac{\partial^2 u}{\partial t^2} = A(x, y, z, t) \frac{\partial^2 u}{\partial x^2} + B(x, y, z, t) \frac{\partial^2 u}{\partial y^2} + C(x, y, z, t) \frac{\partial^2 u}{\partial z^2} + D(x, y, z, t). \quad (1)$$

To solve this equation by the variational iteration method, we need to specify the initial or boundary conditions, which for example may be specified as follows:

$$\text{Initial conditions: } u(x, y, z, 0) = f(x, y, z), \quad \frac{\partial u}{\partial t}(x, y, z, 0) = g(x, y, z).$$

$$\text{Boundary conditions: } u(0, y, z, t) = f_1(x, y, z), \quad \frac{\partial u}{\partial x}(0, y, z, t) = g_1(y, z, t),$$

$$\frac{\partial u}{\partial x}(0, y, z, t) = g_2(y, z, t).$$

For solving Eq. (1) by the iterative variation method, its correction functional can be written down as

$$\text{Case 1: } u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_0^t \lambda_1 \left( \frac{\partial^2 u_n}{\partial \xi^2} - A \frac{\partial^2 \tilde{u}_n}{\partial x^2} - B \frac{\partial^2 \tilde{u}_n}{\partial y^2} - C \frac{\partial^2 \tilde{u}_n}{\partial z^2} - D \right) d\xi$$

$$\text{Case 2: } u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_0^x \lambda_2 \left( \frac{\partial^2 \tilde{u}_n}{\partial t^2} - A \frac{\partial^2 u_n}{\partial \xi^2} - B \frac{\partial^2 \tilde{u}_n}{\partial y^2} - C \frac{\partial^2 \tilde{u}_n}{\partial z^2} - D \right) d\xi$$

$$\text{Case 3: } u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_L^x \lambda_3 \left( \frac{\partial^2 \tilde{u}_n}{\partial t^2} - A \frac{\partial^2 u_n}{\partial \xi^2} - B \frac{\partial^2 \tilde{u}_n}{\partial y^2} - C \frac{\partial^2 \tilde{u}_n}{\partial z^2} - D \right) d\xi.$$

Case 1: To make this correction functional stationary, notice that  $\delta u_n(x, y, z, 0) = 0$ ,

$$\delta u_{n+1} = \delta u_n + (\lambda_1 (\delta u_n)')_0^t - (\lambda_1' \delta u_n)_0^t + \int_0^t \lambda_1'' \delta u_n d\xi = 0.$$

Its stationarity conditions can be obtained as follows:

$$\delta u_n : 1 - \lambda_1'(t) = 0,$$

$$\delta u_n' : \lambda_1(t) = 0,$$

$$\delta u_n : \lambda_1''(\xi) = 0.$$

From this the Lagrange multiplier can be identified as  $\lambda_1 = \xi - t$ , and the following iteration formula will be obtained:

$$u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_0^t (\xi - t) \left( \frac{\partial^2 u_n}{\partial \xi^2} - A \frac{\partial^2 u_n}{\partial x^2} - B \frac{\partial^2 u_n}{\partial y^2} - C \frac{\partial^2 u_n}{\partial z^2} - D \right) d\xi. \quad (2)$$

Starting with  $u_0$ , which can be chosen arbitrarily, Eq. (2) will be used to derive iterative approximations to  $u$ . In this case, considering initial conditions, we use  $u_0$  as  $u_0(x, y, z, t) = f(x, y, z) + g(x, y, z)t$ .

Case 2: To make this correction functional stationary, consider  $\delta u_n(0, y, z, t) = 0$ ,

$$\delta u_{n+1} = \delta u_n - (\lambda_2 A (\delta u_n)')_0^x + ((\lambda_2 A)' \delta u_n)_0^x - \int_0^x (\lambda_2 A)'' \delta u_n d\xi = 0.$$

Its stationarity conditions can be obtained as follows:

$$\delta u_n : 1 + (\lambda(x) A(x, y, z, t))' = 0,$$

$$\delta u_n' : \lambda(x) A(x, y, z, t) = 0,$$

$$\delta u_n : (\lambda(\xi) A(\xi, y, z, t))'' = 0.$$

From this the Lagrange multiplier can be identified as  $\lambda_2 = \frac{1}{A(\xi, y, z, t)}(x - \xi)$ , and the following iteration formula will be obtained:

$$u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_0^x \frac{1}{A(\xi, y, z, t)}(x - \xi) \left( \frac{\partial^2 u_n}{\partial t^2} - A \frac{\partial^2 u_n}{\partial \xi^2} - B \frac{\partial^2 u_n}{\partial y^2} - C \frac{\partial^2 u_n}{\partial z^2} - D \right) d\xi. \quad (3)$$

In this case, considering boundary conditions, we use  $u_0$  as  $u_0(x, y, z, t) = f_1(y, z, t) + g_1(y, z, t)x$ .

In case 3, like for case 2, notice that  $\delta u_n(L, y, z, t) = 0$ ; we obtain  $\lambda_3 = \frac{1}{A(\xi, y, z, t)}(x - \xi)$ , and the following iterative formula will be obtained:

$$u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_L^x \frac{1}{A(\xi, y, z, t)}(x - \xi) \left( \frac{\partial^2 u_n}{\partial t^2} - A \frac{\partial^2 u_n}{\partial \xi^2} - B \frac{\partial^2 u_n}{\partial y^2} - C \frac{\partial^2 u_n}{\partial z^2} - D \right) d\xi. \quad (4)$$

Considering boundary conditions, we use  $u_0$  as  $u_0(x, y, z, t) = f_1(y, z, t) + g_1(y, z, t)(x - L)$ .

**Example 1.** Consider the following equation with the following initial conditions:

$$\begin{cases} \frac{\partial^2 u}{\partial t^2} = A \frac{\partial^2 u}{\partial x^2} + B \frac{\partial^2 u}{\partial y^2} + C \frac{\partial^2 u}{\partial z^2} + D \\ u(x, y, z, 0) = a_1 x + a_2 x^2 + b_1 y + b_2 y^2 + c_1 z + c_2 z^2 \\ \frac{\partial u}{\partial t}(x, y, z, 0) = a'_1 x + a'_2 x^2 + b'_1 y + b'_2 y^2 + c'_1 z + c'_2 z^2 \end{cases}$$

where  $A, B, C$  and  $D$  are constant.

From (2), we have

$$u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_0^t (\xi - t) \left( \frac{\partial^2 u_n}{\partial \xi^2} - A \frac{\partial^2 u_n}{\partial x^2} - B \frac{\partial^2 u_n}{\partial y^2} - C \frac{\partial^2 u_n}{\partial z^2} - D \right) d\xi. \quad (5)$$

Starting with

$$u_0(x, y, z, t) = a_1 x + a_2 x^2 + b_1 y + b_2 y^2 + c_1 z + c_2 z^2 + (a'_1 x + a'_2 x^2 + b'_1 y + b'_2 y^2 + c'_1 z + c'_2 z^2)t$$

by iteration formula (5), we have

$$u_1(x, y, z, t) = a_1 x + a_2 x^2 + b_1 y + b_2 y^2 + c_1 z + c_2 z^2 + (a'_1 x + a'_2 x^2 + b'_1 y + b'_2 y^2 + c'_1 z + c'_2 z^2)t + \left( Aa_2 + Bb_2 + Cc_2 + \frac{1}{2}D \right) t^2 + (Aa'_2 + Bb'_2 + Cc'_2) \frac{t^3}{3},$$

which is an exact solution.

**Example 2.** Let us solve the following partial differential equations:

$$\begin{cases} \frac{\partial^2 u}{\partial t^2} = \frac{1}{6} \left( x^2 \frac{\partial^2 u}{\partial x^2} + y^2 \frac{\partial^2 u}{\partial y^2} + z^2 \frac{\partial^2 u}{\partial z^2} \right) \\ u(x, y, z, 0) = x^2 y^2 z^2 \\ \frac{\partial u}{\partial t}(x, y, z, 0) = -x^2 y^2 z^2. \end{cases}$$

From (2) the variational iteration formula can be constructed as

$$u_{n+1}(x, y, z, t) = u_n(x, y, z, t) - \int_0^t (\xi - t) \left( \frac{\partial^2 u_n}{\partial \xi^2} - \frac{1}{6} \left( x^2 \frac{\partial^2 u_n}{\partial x^2} + y^2 \frac{\partial^2 u_n}{\partial y^2} + z^2 \frac{\partial^2 u_n}{\partial z^2} \right) \right) d\xi.$$

By imposing  $u_0(x, y, z, t) = x^2 y^2 z^2 - x^2 y^2 z^2 t$ , we have the following approximate solutions:

$$\begin{aligned} u_1(x, y, z, t) &= u_0 + x^2 y^2 z^2 \left( \frac{t^2}{2!} - \frac{t^3}{3!} \right) \\ u_2(x, y, z, t) &= u_0 + u_1 + x^2 y^2 z^2 \left( \frac{t^4}{4!} - \frac{t^5}{5!} \right) \\ u_n(x, y, z, t) &= u_0 + u_1 + \cdots + u_{n-1} + \left( \frac{t^{2n}}{(2n)!} - \frac{t^{2n+1}}{(2n+1)!} \right). \end{aligned}$$

Thus we have  $u(x, y, z, t) = \lim_{n \rightarrow \infty} u_n(x, y, z, t) = x^2 y^2 z^2 \sum_{n=0}^{\infty} (-1)^n \frac{t^n}{n!} = x^2 y^2 z^2 e^{-t}$ , which is an exact solution.

**Example 3.** Consider the following partial differential equation with specified initial conditions:

$$\begin{cases} \frac{\partial^2 u}{\partial t^2} = e^{-xt} \frac{\partial^2 u}{\partial x^2} + e^{-yt} \frac{\partial^2 u}{\partial y^2} + e^{-zt} \frac{\partial^2 u}{\partial z^2} \\ u(x, y, z, 0) = x^2 + y^2 + z^2 \\ \frac{\partial u}{\partial t}(x, y, z, 0) = 0. \end{cases}$$

The variational iteration method consists of the following scheme:

$$u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_0^t (\xi - t) \left( \frac{\partial^2 u}{\partial \xi^2} - e^{-x\xi} \frac{\partial^2 u}{\partial x^2} + e^{-y\xi} \frac{\partial^2 u}{\partial y^2} + e^{-z\xi} \frac{\partial^2 u}{\partial z^2} \right) d\xi.$$

Taking  $u_0(x, y, z, t) = x^2 + y^2 + z^2$ , we have

$$\begin{aligned} u_1(x, y, z, t) &= x^2 + y^2 + z^2 \\ &\quad + \frac{2 \left( (e^{-xt}(1+x) - 1) y^2 z^2 + (e^{-yt}(1+y) - 1) x^2 z^2 + (e^{-zt}(1+z) - 1) y^2 x^2 \right)}{x^2 y^2 z^2}. \end{aligned}$$

By using  $u_1$  to compute  $u_2$ , and considering  $u \approx u_2$ , we derive

$$\begin{aligned} u(x, y, z, t) &\approx \frac{1}{y^6 x^6 z^6} (-2t z^6 y^6 x^5 - 24e^{(-yt)} z^6 x^6 y + 2z^5 y^6 x^6 + 2e^{(-2xt)} t^2 y^6 x^2 z^6 + 24e^{(-2zt)} y^6 x^6 z \\ &\quad + 24e^{(-2xt)} y^6 x z^6 + 24te^{(-zt)} y^6 x^6 z - 40e^{(-zt)} y^6 x^6 - 2tz^5 y^6 x^6 + 40e^{(-2yt)} z^6 x^6 \\ &\quad + 40e^{(-2xt)} z^6 y^6 + 12te^{(-2zt)} y^6 x^6 z^2 + 12te^{(-2xt)} y^6 z^6 x^2 + 16te^{(-2xt)} y^6 z^6 x \\ &\quad + 16te^{(-2zt)} y^6 x^6 z + 12te^{(-2yt)} y^2 z^6 x^6 + y^8 z^6 x^6 + z^8 y^6 x^6 + x^8 z^6 y^6 \\ &\quad + 16te^{(-2yt)} z^6 x^6 y + 24e^{(-2yt)} z^6 x^6 y - 6ty^6 z^6 x^6 + 2y^5 z^6 x^6 + y^6 z^6 x^6 + 12te^{(-zt)} z^2 x^6 y^2 \\ &\quad - 2ty^5 z^6 x^6 + 24te^{(-xt)} z^6 y^6 x + 40e^{(-2zt)} x^6 y^6 + 12te^{(-xt)} z^6 y^6 x^2 + 2e^{(-2yt)} t^2 z^6 y^2 x^6 \\ &\quad + 2e^{(-2yt)} t^2 z^6 y^3 x^6 - 40e^{(-xt)} z^6 y^6 + 2z^6 y^6 x^5 - 40e^{(-yt)} z^6 x^6 - 24e^{(-xt)} z^6 y^6 x \\ &\quad - 24e^{(-zt)} x^6 y^6 z + 12te^{(-yt)} z^6 y^2 x^6 + 24te^{(-yt)} z^6 y x^6 - 6t^2 e^{(-zt)} z^2 y^6 x^6 \end{aligned}$$

$$-6t^2 e^{(-xt)} z^6 y^6 x^2 - 6t^2 e^{(-yt)} z^3 y^6 x^6 + 6t^2 e^{(-yt)} z^6 y^6 x^6 + t^2 e^{(-zt)} z^6 y^6 x^6 \\ + t^2 e^{(-yt)} z^6 y^6 x^6 + 2t^2 e^{(-2zt)} z^3 y^6 x^6 + 2t^2 e^{(-2zt)} z^2 y^6 x^6 + 2t^2 e^{(-2xt)} z^6 y^6 x^3).$$

It is worth pointing out that the results in Examples 1–3 are exactly the same as the results from applying the Adomian decomposition method [23].

**Example 4.** Consider the following equation with the following boundary conditions:

$$\begin{cases} \frac{\partial^2 u}{\partial t^2} = A \frac{\partial^2 u}{\partial x^2} + B \frac{\partial^2 u}{\partial y^2} + C \frac{\partial^2 u}{\partial z^2} + D \\ u(0, y, z, t) = a_1 y + a_2 y^2 + b_1 z + b_2 z^2 \\ \frac{\partial u}{\partial x}(0, y, z, t) = a'_1 y + a'_2 y^2 + b'_1 z + b'_2 z^2 \end{cases}$$

where  $A$ ,  $B$ ,  $C$  and  $D$  are constant.

From (3), we have

$$u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \frac{1}{A} \int_0^x (x - \xi) \left( \frac{\partial^2 u_n}{\partial t^2} - A \frac{\partial^2 u_n}{\partial \xi^2} - B \frac{\partial^2 u_n}{\partial y^2} - C \frac{\partial^2 u_n}{\partial z^2} - D \right) d\xi. \quad (6)$$

Beginning with  $u_0(x, y, z, t) = a_1 y + a_2 y^2 + b_1 z + b_2 z^2 + (a'_1 y + a'_2 y^2 + b'_1 z + b'_2 z^2)x$ , by iteration formula (6), we have

$$u_1(x, y, z, t) = a_1 y + a_2 y^2 + b_1 z + b_2 z^2 + (a'_1 y + a'_2 y^2 + b'_1 z + b'_2 z^2)x \\ - \frac{1}{A} \left( \left( a_2 B + b_2 C + \frac{1}{2} D \right) x^2 + (a'_2 B + b'_2 C) \frac{x^3}{3} \right),$$

which again is an exact solution.

**Example 5.** In this example we consider the following partial differential equations:

$$\begin{cases} \frac{\partial^2 u}{\partial t^2} = x \frac{\partial^2 u}{\partial x^2} + y \frac{\partial^2 u}{\partial y^2} + z \frac{\partial^2 u}{\partial z^2} \\ u(1, y, z, t) = yzt^2 \\ \frac{\partial u}{\partial x}(1, y, z, t) = -yzt^2. \end{cases}$$

From (4) the variational iteration formula will be obtained as

$$u_{n+1}(x, y, z, t) = u_n(x, y, z, t) + \int_1^x \frac{1}{\xi} (x - \xi) \left( \frac{\partial^2 u_n}{\partial t^2} - \xi \frac{\partial^2 u_n}{\partial \xi^2} + y \frac{\partial^2 u_n}{\partial y^2} + z \frac{\partial^2 u_n}{\partial z^2} \right) d\xi.$$

By imposing  $u_0(x, y, z, t) = 2yzt^2 - xyz t^2$  we have

$$u_1(x, y, z, t) = 2yzt^2 + 4xyz \ln x - x^2 yz - 2xyz + 3yz,$$

which is an exact solution.

### 3. Conclusion

In this work, we present the analytical approximation to a solution for wave equations in three different cases. We have achieved this goal by applying He's variational iteration method. Using the variational iteration method, it is possible to find the exact solution or a good approximate solution of the equation. It can be concluded that He's variational iteration method is very powerful and efficient technique for finding exact solutions for wide classes of problems. One of the restrictions in this method is the selection of  $u_0$  relative to initial and boundary conditions. The suitable selection of  $u_0$  has a very significant effect in computing  $u$ . Computations are performed using the Maple 10 package.

## References

- [1] J.H. He, Variational iteration method—some recent results and new interpretations, *J. Comput. Appl. Math.* 207 (1) (2007) 3–17.
- [2] J.H. He, Variational iteration method for autonomous ordinary differential systems, *Appl. Math. Comput.* 114 (2–3) (2000) 115–123.
- [3] J.H. He, X.H. Wu, Construction of solitary solution and compaction-like solution by variational iteration method, *Chaos Solitons Fractals* 29 (2006) 108–113.
- [4] Shu-Qiang Wang, Ji-Huan He, Variational iteration method for solving integro-differential equations, *Phys. Lett. A* 367 (3) (2007) 188–191.
- [5] J.H. He, Variational iteration method—a kind of nonlinear analytical technique: Some examples, *Internat. J. Non-linear. Mech.* 34 (1999) 708–709.
- [6] J.H. He, Some asymptotic methods for strongly nonlinear equations, *Internat. J. Modern Phys. B* 20 (10) (2006) 1141–1199.
- [7] J.H. He, Approximate analytical solution for seepage flow with fractional derivatives in porous media, *Comput. Methods Appl. Mech. Engrg.* 167 (1998) 57–68.
- [8] J.H. He, A new approach to nonlinear partial differential equations, *Commun. Nonlinear Sci. Numer. Simul.* 2 (4) (1997) 203–205.
- [9] J.H. He, A variational iteration approach to nonlinear problems and its applications, *Mech. Appl.* 20 (1) (1998) 30–31.
- [10] J.H. He, Variational iteration method for delay differential equations, *Commun. Nonlinear. Sci. Numer. Simul.* 2 (4) (2007) 235–236.
- [11] S. Momani, S. Abuasad, Application of He's variational iteration method to Helmholtz equation, *Chaos Solitons Fractals* 27 (5) (2005) 1119–1123.
- [12] M.A. Abdou, A.A. Soliman, Variational iteration method for solving Burgers' and coupled Burgers' equations, *J. Comput. Appl. Math.* 18 (2005) 1245–1251.
- [13] M. Tatari, M. Dehghan, He's variational iteration method for computing a control parameter in a semi-linear inverse parabolic equation, *Chaos Solitons Fractals* 33 (2) (2007) 671–677.
- [14] Abdul-Majid Wazwaz, A study on linear and nonlinear Schrodinger equations by the variational iteration method, *Chaos Solitons Fractals* (in press).
- [15] J. Biazar, H. Ghazvini, He's variational iteration method for solving hyperbolic differential equations, *Int. J. Nonlinear Sci. Numer. Simul.* 8 (3) (2007) 311–314.
- [16] J. Biazar, H. Ghazvini, He's variational iteration method for fourth-order parabolic equations, *Comput. Math. Appl.* (in press).
- [17] J. Biazar, H. Ghazvini, He's variational iteration method for solving linear and non-linear system of ordinary differential equations, *Appl. Math. Comput.* 191 (1) (2007) 287–297.
- [18] E. Yusufoglu, Variational iteration method for construction of some compact and noncompact structures of Klein–Gordon equations, *Int. J. Nonlinear Sci. Numer. Simul.* 8 (2) (2007) 152–158.
- [19] Z.M. Odibat, S. Momani, Application of variational iteration method to Nonlinear differential equations of fractional order, *Int. J. Nonlinear Sci. Numer. Simul.* 7 (2006) 27–34.
- [20] N. Bildik, A. Konuralp, The use of variational iteration method, differential transform method and adomian decomposition method for solving different types of nonlinear partial differential equations, *Int. J. Nonlinear Sci. Numer. Simul.* 7 (2006) 65–70.
- [21] E.M. Abulwafa, M.A. Abdou, A.A. Mahmoud, The solution of nonlinear coagulation problem with mass loss, *Chaos Solitons Fractals* 29 (2006) 313–330.
- [22] S. Momani, Z. Odibat, Analytical approach to linear fractional partial differential equations arising in fluid mechanics, *Phys. Lett. A* 1 (53) (2006) 1–9.
- [23] J. Biazar, R. Islam, solution of wave equation by Adomian decomposition method and the restrictions of the method, *Appl. Math. Comput.* 149 (2004) 807–814.